






Thank you for attention...



YSOs
- A rich number of Young Stellar Objects observed with VEGA
- Study of the S1 Gps
N. Jamshidshahi, K. Fomart, M. Bantay, M. Bani, A. Wafar, R. Lopez, P. et al., D. Mousavi, C. Desgagnés, G. Lenc



Observatoire de la cote d'Azur

Narges Jamialahmadi

Supervisors:

Bruno Lopez; philippe Berio





YSOs

- *A mini review of Young Stellar Objects observed with VEGA*
- *Study of the 51 Oph*

*N. Jamialahmadi; K. Perraut; M. Benisty; Ph. Berio
A. Meilland; B. Lopez; Ph. stee;
D. Mourard; C. Dougados; G. Lima;*



Scientific rationale

- *Understand the physical mechanism involved around YSOs...*
- *Spatially and spectrally resolved observations of such targets across the H α line with VEGA is crucial as it will enable the relative contributions of the accretion and ejection processes to the line formation to be disentangled.*
- *Bring direct spatial constraints on the geometry of the winds, and on the accretion/ejection scenario at scales as small as 0.1 AU.*


Target	Sp. type	L (L_{\odot})	R_{sub} (AU)	d (pc)	R_{sub} (mas)	Ha line
MWC 361	B2Ve	3000 – 15000	3.7 – 8.4	360	10 - 23	Single- or double-peaked Benisty, Perraut et al. 2013, A&A, 555, A113
AB Aur	A0Ve	~ 50	0.5	144	3.4	(variable) P Cygni profile Perraut et al. 2010, A&A, 516, L1 Lima, Perraut et al. in prep
MWC 275	A1Ve	~ 30000	11.9	122	97	(variable) P Cygni profile
MWC 158	B6V[e]	~ 10000	6.9	500	14	Double-peaked Ellerbroek, Benisty et al. in prep.
MWC 480	A3Ve	~ 10	0.2	170	1.3	(variable) P Cygni profile
51 Oph	B9.5IIIe	~ 260	0.54	131	4	Double-peaked Dunkin et al. 1999




MWC361: enhanced Ha activity at periastron

[Benisty, Perraut et al. A113] 2013, A&A, 555

3-year follow-up of the young binary system MWC361





VEGA Ha line over the orbital period



VEGA differential visibilities (S1S2) across the Ha line (observations and best model)

MWC361: an enhanced mass-loss event

[Benisty, Perraut et al. 2013, A&A, 555, A113]




outburst of accretion, followed by a massive ejection, at the periastron

AB Aur: temporal variability

[Lima, Perraut et al, A&A, in prep.]

3-year follow-up VEGA observations:




20 Ha VEGA spectra exhibiting a P-Cygni profile and a fast variability mainly in the blue wing

AB Aur: constraining the wind

[Lima et al. 2010, A&A, 522, 104]

Simultaneous fit of (variable) spectra and interferometric data allow to:

- constrain the disk wind
- study of magnetospheric accretion



Four components:

- the star (1)
- the magnetosphere (2)
- the accretion disk (3)
- the disk wind (4)





AB Aur: constraining the wind

$i = 30^\circ$

Disk wind: $R_{in} = 5 R^*$, $R_{out} = 25 R^*$, launching angle = 44° , $T_{eff} = 7500 K$

Mass loss rate = $1.7e-08 M_{sun}/yr$ without magneto


[Lima, Perraut et al, A&A, in prep.]



HD50138: resolving the gas disk

Pre- or post-main sequence B[e] star HD50138


[Ellerbroek, Benisty et al, A&A, in prep.]



VEGA differential visibilities (S1S2) across the H α line

emission size of $\sim 1-3$ mas more consistent with an outflow than accretion

VEGA data included in a **multi-technique, multi-lambda** study (X-shooter spectra, AMBER K-band and Brgamma data, PIONIER H-band)




IR interferometric data



51 Oph presentation



<u>Alternate name</u>	<u>Spectral type</u>	<u>Distance (pc)</u>	<u>Velocity (v sini) (km/s)</u>	<u>Magnitude (V)</u>	<u>Age (Myr)</u>	<u>Mass (x stellar mass)</u>
<i>HD 158643</i>	<i>B9.5IIIe</i>	<i>131</i>	<i>267 ± 5</i>	<i>4.83</i>	<i>0.3</i>	<i>4</i>



Evolutionary status of 51 Oph ???

Be star???

Herbig Ae/Be star???



Beta Pictoris???



- 1) Emission lines
- 2) Infrared excess
- 10 micron silicate feature
- Presence of circumstellar dust



- 1) Fast-rotating
- 2) Hot molecular lines emission
- 3) The compact gaseous disk



- 1) Edge-on disk with both gas and dust
- 2) Variable absorption features suggesting infalling gas and materials
- 3) Rare nearby example of a young debris disk with gas just entering the late stages of formation



- 1) Hot CO emission
- 2) Lack of near infrared excess



IR emission interpreted as due to free-free emission from an ionized high-density envelope around star




- 1) Lacking the far infrared-excess bump (No dust detection at 18 μm)
- 2) Absence of H alpha emission in Beta Pic
- 3) Beta Pic star age of 8-12 $\times 10$ Myr and mass of 1.75 stellar mass

VEGA Observations:

May (2013) → E1E2W2; Run: VEGA MR: In continuum

July (2013) → S1S2; Run : VEGA MR : H alpha (two points)

→ W1W2; Run : VEGA MR: poor quality





→ Fundamental parameters of this star

→ Study of the gaseous disk



Data process in continuum neighbored Halpha

Results of first processing:



From data obtained in May 2013,

**Estimated angular diameter:
0.39 +/- 0.01 mas**

**Assuming all flux in the continuum comes
from the central star**





Calculating the effective temperature:



First estimation


$$T_{eff} \sim \left(\frac{L_{\star}}{4\pi\theta_{LD}^2 d^2 R_{\odot}^2 \sigma} \right)^{1/4}$$



T_eff= 9500 +62/-53 K

Estimated with angular diameter in one direction...

**L (star)= 260 +60/-50L (sun) ; (Thi et al . 2013)
d=131 +/- 4 pc (Perryman et al. 1997)**













Results of second processing:

Processing for:



data obtained in May (baseline E1E2; 1 point)

data obtained in July (baseline S1S2; 2 points)



LESIA



Observatoire de Toulouse

Max-Planck-Institut für Radioastronomie






Are VEGA data consistent with a keplerian disk?

We use: Keplerian and uniform disk model

Free parameters:

- 1) Inclination**
- 2) P.A.**
- 3) Major-axis FWHM in the continuum**
- 4) Major-axis FWHM in the line**





Derived information:

For star:


- 1) The angular diameter= 0.39 ± 0.01 mas
- 2) First estimation of the effective temperature= $9500^{+62/-53}$ K

For gaseous disk:

- 1) *Gaseous disk is keplereian* \rightarrow According with Thi et al. 2005
- 2) $i=88$ \rightarrow According to (Gil et al . 2006) and (Tatulli et al. 2008)
- 3) P.A. = 197 \times In Tatulli et al 2008 \rightarrow P.A=129
- 4) FWHM in the continuum= 3 stellar diameter
- 5) FWHM in the line= 10 stellar diameter



New observations 2014:



First run: E2S2W2 or W1W2E2 or W1W2S2;
VEGA MR : Continuum



To constrain the photosphere
(extension and flattening) of
this close-to-critically rotating
star

Second run: VEGA MR: Halpha S1S2; E1E2;
W1W2

To improve our knowledge on the gaseous disk
geometry and kinematics and confirm the estimated
parameters of the keplerian disk





Conclusions:

For YSOs studies:

- 1) Importance of adding the IR data in muliti wavelenghts to visibile interferometric data
- 2) New constrains to modelize the physics involved (wind, accretion, magnetospheric) models.

Thank you for attention...